

Videonystagmography (VNG) Medication Discontinuation List

Do not take any of the following medications for 48 hours prior to your VNG test date. **If you have questions or concerns about stopping any of your medications, please contact your physician.**

Anti-Dizziness Medications:

- Antivert, Bonine (meclizine)
- Benadryl (diphenhydramine)
- Dramamine (dimenhydrinate)

Sedatives:

- Transderm Scop patches
- Vontrol

Antinausea Medications:

- Phenergan

Antibiotics Medications:

- Gentamicin
- Streptomycin

Anticonvulsants medications:

- Klonopin (do not stop for anxiety disorder)
- Dilantin (phenytoin), Lirium, Librax
- Tegretol (carbamazepine), Nembutal

Antidepressants Medications:

- Antivan
- Elavil
- Lithium
- Pamelor
- Placidyl
- Prozac
- Valium
- Xanax
- Cymbalta

Antihistamine/Decongestion Medications:

- Allegra
- Benadryl
- Chlor-Trimeton
- Claritin
- Seldane
- Sudafed
- Zyrtec

Sleeping Medications:

- Ambien
- Dalmane
- Doral
- Halcion
- ProSom
- Restoril
- Seconal
- Unisom

Stimulant Medications:

- Amphetamine
- Ritalin

Headache Medications:

- Fiorinal
- Fioricet

Other Medications:

- Methadone
- NSAID
- Quinine
- Salicylates-Aspirin